

*Reference  
made in  
pleural  
mesothelioma*

PLAINTIFF'S  
EXHIBIT  
P-5-40 ID

## DIFFUSE PLEURAL MESOTHELIOMA AND ASBESTOS EXPOSURE IN THE NORTH WESTERN CAPE PROVINCE

BY

J. C. WAGNER, C. A. SLEGGs, and PAUL MARCHAND

*From the Pathology Division, Pneumoconiosis Research Unit of the Council for Scientific and Industrial Research, Johannesburg, West End Hospital, Kimberley, and the Department of Thoracic Surgery, University of the Witwatersrand and Johannesburg General Hospital*

(RECEIVED FOR PUBLICATION APRIL 24, 1960)

Primary malignant tumours of the pleura are uncommon. Thirty-three cases (22 males, 11 females, ages 31 to 68) of diffuse pleural mesothelioma are described; all but one have a probable exposure to crocidolite asbestos (Cape blue). In a majority this exposure was in the Asbestos Hills which lie to the west of Kimberley in the north west of Cape Province. The tumour is rarely seen elsewhere in South Africa.

Mesothelioma of the pleura is regarded as an uncommon tumour. In the last four years we have seen 33 histologically proven cases; 28 of these had some association with the Cape asbestos field and four cases had been exposed to asbestos in industry.

The tumour is rarely encountered elsewhere in South Africa. During the past five years, with the exception of the present series, no neoplasm of this nature has been diagnosed amongst 10,000 lungs examined at the Pneumoconiosis Bureau in Johannesburg, or in the Pathology Department of the South African Institute for Medical Research. Higginson and Oettle (1957) did not observe a single case in their survey of malignant tumours occurring in the Bantu and Cape Coloured population of Johannesburg and the North Eastern Transvaal.

Our first necropsy specimen of pleural mesothelioma with asbestosis was examined at the Pneumoconiosis Research Unit in February, 1956 (Case 1). During the early months of that year, one of us (C.A.S.) in the Northern Cape, treated six patients with gross pleural thickening. Pleural biopsies from two of them showed the features of mesothelioma. In the ensuing two years, eight further cases were found from this region and five from elsewhere in the Union. During this period C.A.S. had become perturbed at the number of these unusual tumours occurring amongst his patients, and stimulated an investigation. At this stage there were two reasons to suggest that asbestos might be implicated. First, asbestos was found in

the lungs of the first case (Case 1), and secondly 10 of the cases came from a hospital to which suspected cases of tuberculosis were referred from large asbestos mining area. This hypothesis could not be supported at once from the original histories obtained from the patients, for they included housewives, domestic servants, cattle herders, farmers, water bailiff, an insurance agent, and an accountant none of whom were working on the asbestos mines at the time. We therefore undertook a detailed investigation of their past occupation and place of residence, and the association with asbestos exposure was discovered. The cases are summarized in Table 3. Previous biopsy specimens were re-examined, and new cases diagnosed, including asbestos miners. In only one case do the relatives deny that the patient either visited the asbestos mines or was exposed to asbestos.

This is a preliminary publication and the problem is being intensively investigated.

### The Asbestos Area of the North-West Cape

According to Hall (1930) asbestos was discovered by Lichenstein near Prieska during his travels between 1803 and 1806. Since then it has been established that the asbestos deposits extend from 20 miles south of Prieska, northwards through the western part of the magisterial district of Hay, to the eastern portion of the magisterial district of Postmasburg and finally, to the western area of the




FIG. 1.—Map of Griqualand West asbestos fields.

district of Kuruman (Fig. 1). These deposits occur in the slopes of a range of hills covering an area of approximately 8,000 square miles. Known as the "Asbestos Mountains", these hills extend more or less longitudinally between 22°30' and 23° E. The whole area is semi-arid, sparsely populated, and, apart from mining, cattle ranching is the only important occupation. In the Kuruman area several large tracts of land in or abutting on these mountains have been reserved for the aboriginal inhabitants and here whites may only reside in the immediate vicinity of the mines.

The type of asbestos mined throughout this area is crocidolite, better known as Cape Blue Asbestos. The chemical analysis of the fibre is given in Table I. Crocidolite is the fibrous form of riebeckite and all stages of transformation from a massive riebeckite rock through lamellar riebeckite to asbestiform crocidolite occur in this region. Magnetite is frequently associated with the crocidolite (Vermaas, 1952).

Mining of asbestos first began in the Prieska district in 1893 and gradually spread northwards.

In 1908 production had begun in the Kuruman district. Between 1916 and 1918 a large number of claims were taken up. This northward trend has continued and in about 1950 mining started at Pomfret near the Bechuanaland border (Table 2).

Initially the ore was quarried in numerous small open cast workings. This was followed gradually

TABLE I  
CHEMICAL ANALYSES OF CROCIDOLITE AND AMOSITE  
(from Vermaas, 1952).

	Crocidolite	Amosite
S <sub>2</sub> O <sub>3</sub>	31.94	49.47
Al <sub>2</sub> O <sub>3</sub>	0.20	0.63
Fe <sub>2</sub> O <sub>3</sub>	18.64	4.15
FeO	19.39	35.43
MgO	1.37	9.37
CaO	0.19	0.52
Na <sub>2</sub> O	6.07	0.02
K <sub>2</sub> O	0.04	0.20
H <sub>2</sub> O	0.31	0.07
H <sub>2</sub> O+	2.58	2.33
TiO <sub>2</sub>	—	0.25
MnO	—	0.61
F	—	0.01
Total	100.73	100.46

TABLE 2  
POPULATION OF THE MAGISTERIAL DISTRICTS IN WHICH THE ASBESTOS FIELDS ARE SITUATED  
(taken from Union Census 1921, and 1931)

District	Population 1921			Population 1931		
	Whites	Non-whites	Total	Whites	Non-whites	Total
Prieska	3,430	4,879	8,309	3,361	10,847	14,208
Kuruman	4,713	16,498	21,211	4,818	23,779	28,597
Ilav	4,499	8,814	13,313	2,757	9,631	12,388
Postmasburg		Included in the Hay District		4,887	21,019	25,906

by a type of shallow mining; inclined shafting became more common after 1930.

According to Frood (1915), the quarrying was a family affair undertaken by the local inhabitants. The men quarried the rock, which was sorted and then hand cobbled by the women and children. Hand cobbing consists of separating the fibre from the banded ironstone by striking the rock cob with a small hammer. As the market was selective, a rotary sieve was sometimes used to grade the fibre and eliminate the small particles of dust. After grading, it was bagged and weighed. The asbestos was then transported by donkey wagon to the nearest rail-head. Before the establishment of the railway in this area between 1923 and 1930, such a journey could last as long as 10 days. This transport was generally undertaken by white youths (farmers' sons) under contract. The white inhabitants played very little part in actual recovery of the fibre, being nearly always employed in the role of managers or overseers, issuing stores and supervising the grading and weighing of the fibres. Farming activities were and are still carried out round the mines.

Since the 1939-1945 war, the demand for crocidolite has enormously increased. The lucrative claims have been bought by registered companies, others have been abandoned. As deeper and richer deposits have been found, vertical shafts are being sunk. However, the mines with large shallow deposits are still using the inclined shaft and tunnel, and quarrying is still used by the few remaining smaller producers. With the building of more mills, hand cobbing has diminished. In 1915 the first crushing mill was established at Koegas in the south. This was followed by a large mill at Kuruman (operated between 1926 and 1931) where it was situated within 300 yards of the main street, close to which cobbing was also done for a few years (1927-1930). This was followed by a mill at Prieska in 1930, which was completely rebuilt in 1957. Griquatown had a small mill in 1928. The practice today is for one mill to serve several mines in the immediate vicinity.

In the early days the manager and labourers lived within a few yards of their place of work, and even

today the non-white prefers to live as close as possible, and the children play on the dumps from the mine and mill.

#### Case Histories

The following eight case histories illustrate various aspects of the disease and the different types of exposure to asbestos dust.

Case 1.—B.P., a Bantu male, 36 years of age (born 1920), was a mine labourer, and was the first case diagnosed as a mesothelioma with evidence of asbestos. He was born in the Kuruman district but it is not known whether he worked in the asbestos mines. He was employed on the Witwatersrand gold-mines underground for two years and in the change rooms for a further 11 years.

A radiograph taken at a mine hospital on August 16 1955 showed a massive right-sided pleural effusion and 3,000 ml. of fluid was withdrawn. He was admitted to the Witwatersrand Native Labour Association Hospital on August 24, 1955, and two days later aspiration yielded 1,000 ml. of thick gelatinous pus "which could be pulled out in threads". He was treated with frequent aspirations and instillation of varidase but without improvement, and he died on February 15, 1956 (Martiny, 1956).

At autopsy the right thoracic cavity was occupied by a large gelatinous tumour which displaced the mediastinum and compressed the left lung. The tumour had infiltrated the pericardium. The right lung was completely compressed by neoplastic tissue (Fig. 2) but the right bronchial tree did not show any evidence of primary bronchogenic carcinoma. Histological sections of the pleural growth showed a papillary mesothelioma (Fig. 3). There was evidence of asbestosis in both lungs (Fig. 4).

Case 4.—K.H., a white female, 56 years of age (born 1898), was a social worker, who could only have had a short exposure to asbestos as a child and probably a further slight exposure as a young woman. She may have paid several short visits to the mines with her husband at a later period.

She was born in Griquatown where she lived until she was 5 years of age. Her family then moved to Kimberley where from 1916 to 1922 she worked as a clerk in an asbestos warehouse. Her husband owned an asbestos mine from 1933-1940.



FIG. 2

2.—Right lung of Case 1 showing a tumour compressing the lung substance.



FIG. 3

3.—Cellular structure of tumour of Case 1.  $\times 40$  Haematoxylin and eosin.



FIG. 4

4.—Sections of lung from Case 1 showing asbestos bodies.  $\times 40$  Haematoxylin and eosin.



FIG. 5.—On July 21, 1954.




FIG. 6.—On July 7, 1955.




FIG. 7.—On March 3, 1956.





FIG. 8.—On July 3, 1956.

FIGS. 5, 6, 7, and 8.—Radiographs showing development of tumour in Case 4.



FIG. 9.—Right lung of Case 14.

FIG. 10—Histological sections of tumour from Case 14.  $\times 26$   
Harrimantoxilin and eosin.

he was referred to the West End Hospital, Kimberley, July 19, 1954, because of dyspnoea and right chest discomfort of acute onset. The chest radiograph showed a loculated pleural effusion at the right base. Straw-coloured fluid was aspirated, which on examination was found to be negative for *M. tuberculosis* but stained mesothelial cells arranged in acini. At necropsy in Johannesburg on August 11, 1954, four nodules were seen on the parietal pleura. Histological examination of the biopsy specimen showed a mesothelioma. Deep therapy and radio-active gold implantation in September, 1955 caused initial regression of the pleural fluid and nodules but did not arrest the progress of the disease. She received a further course of radiotherapy without improvement and died in January, 1957. The liver was enlarged and nodular on gross death but no necropsy was done. (Figs. 5, 6, 7, 8 show the radiological development of the tumour in his case.)

Case 14.—S.S., a Nantu male, 40 years of age (born 1888), was an asbestos miner. He was born in the Kuruman district and as a child often played on the asbestos dump near an asbestos mine. He subsequently worked with asbestos for several years, being employed at an asbestos mine weighing fibre from 1938-41. In 1942 he worked on a Witwatersrand gold mine but

returned to the Kuruman district in 1943 where he worked as a farm labourer until his final illness.

He was admitted to the West End Hospital on March 3, 1959. He had been well until July, 1958 when he became aware of pains in the right side of his chest. In addition he had a slightly productive cough and shortness of breath on exertion. Chest radiograph on March 9, 1959 showed obliteration of the right lung field. Asbestos bodies were found in the sputum. Needle biopsy showed histological features consistent with a mesothelioma.

**Autopsy.**—The right chest was occupied by a huge whitish tumour which infiltrated the mediastinum, extending through the diaphragm onto the superior surface of the liver. A straw-coloured effusion was present in the left pleural sac but the left lung was not affected. The mediastinal, cervical, and para-aortic abdominal lymph glands to the level of the renal arteries were infiltrated with tumour.

In the right lung it was found that the tumour mass had infiltrated along the interlobar fissures. The lobes of the lung were compressed into three small segments (Fig. 9). On histological section a papillary type of pleural mesothelioma was seen (Fig. 10) which had infiltrated into the lung parenchyma in some areas. There was marked lymphatic spread throughout the

lung substance. Numerous asbestos bodies were present in the air spaces and occasional bodies and fibres were seen in the interstitial tissue. The lung itself was markedly atelectatic. Evidence of asbestosis was observed in the left lung.

**Case 15.**—S.S.V., a white female, 42 years of age (born 1916), is a housewife, who lived on a mine near Johannesburg and was treated for tuberculous pleurisy for six months. It was only after the histological diagnosis of a mesothelioma had been made that her previous history was obtained. In addition, it was subsequently learnt that her father (Case 22) had been born on the asbestos field and had a long mining history.

The patient was born in the Griquatown district and went to school at Kuruman. She spent most of her youth in the asbestos belt. She has lived near Johannesburg for 20 years.

In March, 1958 this patient developed right chest pain and rapidly increasing shortness of breath. She was admitted to hospital and 2,500 ml. of slightly blood-stained fluid was removed from the right chest. A thorascopic examination was carried out on July 27, when a small biopsy was taken but this consisted of fibrous tissue. Bronchoscopy was normal and no neoplastic cells were observed in the bronchial washings. *M. tuberculosis* was not cultured from numerous specimens of sputa, pleural fluid nor from the bronchial washings. She was admitted to a Johannesburg nursing home on October 16, 1958.

Radiographs then showed massive thickening of the right pleura with fluid in the right chest. At thoracotomy (Mr. Dennis Fuller) the parietal pleura was 2 cm. thick, "hard as rock" and stripped with great difficulty from the chest wall. The visceral peel stripped fairly easily from the upper lobe but separation became more difficult over the lower lobe where the peel was thicker. During the decortication a necrotic mass was found in the parenchyma of the left lower lobe. At the base, confluent with the diaphragm and perhaps the mediastinum, was a hard craggy mass which could not be removed. Histological sections of the specimen revealed an adenoid form of tubular mesothelioma. Subsequently three specimens of sputum were examined for asbestos bodies but none were found.

The patient is still alive (April, 1960), but her condition is deteriorating.

**Case 22.**—D.A.B., a white male, 68 years of age, and a farmer, was the father of the previous patient. Although a diagnosis of pleural mesothelioma was made in January, 1957 it was not until July, 1959 that details of his exposure to asbestos was obtained.

He was born in Griquatown and went to school there. He worked in Kuruman, Posinasburg and Griquatown until the age of 62. Between 1918 and 1932 he worked as an asbestos miner underground and was mostly in charge of blasting. Between 1932 and 1944 he farmed in the district but returned to asbestos mining from 1945 to 1952.

In January, 1957 the patient consulted his doctor because of left-sided chest pain with cough and slight

haemoptysis. He was referred to the Johannesburg General Hospital for investigation. The chest radiograph showed gross pleural thickening with a basal pleural effusion on the left side. He was admitted to hospital and on June 10, 1957 a thorascopic pleural biopsy was taken. This showed the features of a mesothelioma. On June 24, 1957 an attempt at decortication was made. The left upper lobe was stripped easily but the grossly thickened pleura was so adherent to the lower lobe that the attempt was abandoned. Sections of the pleura confirmed the diagnosis of mesothelioma. He died on December 13, 1957.

**Case 16.**—A.N.V., a white male, 50 years of age (born 1906), was a farmer, who was born at Griquatown and went to a village school near an asbestos mine. From 1914-16 he transported asbestos in donkey waggons from Hay to Prieska. During these journeys he often slept in asbestos bags. In 1914 he worked in an asbestos mine for a month.

The patient became ill in December, 1955 with vague substernal pain and dyspnoea. A left-sided pleural effusion was diagnosed and he was referred to Mr. J. Bremmer in Pretoria. There was no evidence of malignancy in bronchial and scalene biopsies. Some apical pleural thickening and a dense homogeneous opacity at the base of the left lung was seen in the radiograph August 3, 1956. Thoracotomy (Mr. Bremmer) February, 1957 revealed a thickened nodular pleura with a large mass on the left diaphragm. A mesothelioma was seen on histological section.

In April, 1959 he was examined by Mr. J. D. Visker Kimberley who found small nodules in the subcutaneous tissue over the neck and chest. A large firm mass was found in the thoracotomy wound and a mass was palpable in the left hypochondrium. There was general lymphadenopathy and numerous nodules were present on the tongue. Chest radiographs showed the left pleural cavity to be obliterated. He was one of the few cases to develop widespread metastases.

A biopsy of one of the subcutaneous chest nodules showed a similar appearance to those seen in the original specimen from the pleura. The patient died on May 1959. No necropsy was performed.

**Case 24.**—D.G., a white male, 32 years of age (born 1922), by profession an accountant, was born in England but came to South Africa as an infant. He spent his early childhood in Kuruman and left at the age of 7. He later qualified as a chartered accountant in Kimberley. In 1944, at the age of 22, he had been discharged medically unfit from the army because of a left pleural effusion. Neither then nor on many subsequent occasions were acid-fast bacilli found, either in sputum or pleural fluid. His only subsequent contact with asbestos was auditing of the books of an amosite mine in Transvaal.

In September, 1953 he complained of left-sided chest pains and shortness of breath. A radiograph of the chest showed flattening of the diaphragm on the left side, obliteration of the costophrenic angle and thickening of the oblique fissure. He was seen on numerous occasions




FIG. 11.—On September 11, 1956.




FIG. 12.—On August 23, 1957.

FIGS. 11 and 12.—Radiographs showing development of tumour in Case 24.

1953 to 1957. A radiograph taken early in 1956 showed the development of a left superior mediastinal shadow which enlarged progressively (Fig. 11) and later development of diffuse nodular pleural thickening on the left. During 1957 similar but less marked changes developed on the right side (Fig. 12). He consented to surgery in June, 1957, and, at operation, a diffuse pleural malignancy was present.

Sections of the parietal pleura only were removed and histological appearances were those of a mesothelioma. He died on October 6, 1957. A necropsy was performed.

Case 30.—G.F., a white male, 50 years of age (born 1887), is a storekeeper, with a history of industrial silture. He has never been near the Cape Asbestos works.

He was born in South West Africa and lived there until the age of 21. From the age of 21 to 26 he worked in a factory in the Cape as a boiler-maker and fitter. He received the information that he had worked continuously with asbestos, lagging pipes and boilers. He remarked that the removal of asbestos lagging from old buildings created a very dusty atmosphere and that he would get a lot of dust at such work. Between 26 and 32 years of age he worked in a factory in Johannesburg in a similar capacity but was not exposed to such heavy dust conditions. For the past 20 years he had not been in contact with asbestos. Enquiries were made from the factory in the Cape as to the nature of the asbestos used

and these have revealed that during the relevant periods Cape crocidolite was exclusively in use.

He was first seen on March 14, 1959 complaining of cough, pain in the right chest and recent onset of breathlessness. Three pints of fluid had been aspirated from the pleura and cytological examination showed the presence of malignant cells. The radiograph showed marked pleural thickening and hydro-pneumothorax. On March 17, 1959, a thoracoscopy was done. The pleura felt thickened and the lung was collapsed and immobile. Several biopsies were taken. These showed the presence of a mesothelioma. The underlying lung showed the presence of asbestosis. He was treated with nitrogen mustard intra-pleurally and his condition temporarily improved. When last examined he complained of increased breathlessness.

#### Discussion

In 1924 Robertson denied the existence of primary malignant tumours of the pleura and considered them to be secondary in origin. Since then, on the one hand, Willis (1948, 1953) and Smart and Hinson (1957) have supported Robertson's views, while on the other hand, primary neoplasms of this nature have been described by many authors in recent years. These include Tobiassen (1955) in Sweden, Belloni and Bovo (1957) in Italy, Godwin (1957) in the United States, and McCaughey (1958)



in Britain. Evidence in support of the mesothelial origin of these tumours can be found in the tissue culture experiments of Stout and Murray (1942) and Sano, Weiss, and Gault (1950).

The variegated histological pattern of tumours arising from the mesothelium of the pleura was remarked on by Klempner and Rabin (1931). Such a variation is appreciated when the multipotentiality of the cells lining coelomic cavities are considered. Maximow first demonstrated this feature in 1927. Novak (1931) stated that the mucosa of all parts of the Müllerian canal and the germinal epithelium were derived from these cells. Keasbey (1947) showed that in the embryo, the mesonephros, metanephros, Wolffian body, genital ridge, and all dependent urogenital structures are derived from the mesothelium.

Campbell (1950) considered the presence of both epithelial and mesenchymal elements a major diagnostic feature. In describing the histology of 11 of these tumours McCaughey (1958) demonstrated that either the epithelial or the mesenchymal element might predominate. He classified his cases into the following four groups: Tumours of epithelial character; tumours of mesenchymal type; tumours of mixed type; and tumours of anaplastic type.

Using McCaughey's classification on our series of tumours, the majority are, as in Campbell's series, of the mixed type. Of the remainder, a few showed the papillary tubular structure of the "tumours of epithelial character", and there were several of the anaplastic type. One case had the appearances of a "tumour of mesenchymal type" but even in this case primitive tubular structures were seen.

Apart from the original case all the histological diagnoses were made on biopsy material. Three of these biopsy findings were confirmed at necropsy. In the remainder, we have had to rely on clinical and radiological examination to exclude other primary sites of malignancy.

Following the work of Meyer and Chaffee (1939, 1940), the possibility of demonstrating hyaluronic acid in these tumours both chemically and histochemically has been considered as a diagnostic aid. Both these investigations are still in an early stage (Harrington, 1959).

Preliminary results of the histochemical experiments show that there is metachromatic substance in both the stroma and glandular structures of the tumours. This metachromasia can be reversed by incubation with testicular hyaluronidase. This material stains strongly with Hale's (1946) colloidal iron method. The periodic acid Schiff technique has given variable reactions. These results are not specific for hyaluronic acid but are strongly suggestive of its presence.

The amount of metachromatic substance in the tumours has varied considerably. This we think partly due to the fact that until recently we had not appreciated the solubility of hyaluronic acid in aqueous media and the majority of these tumours were fixed in 10% formal-saline. Further, Lise (1953) states that the metachromasia of hyaluronic acid is optimal at a concentration of 1/10,000 and gradually decreases at higher levels. When the low concentrations that Meyer and Chaffee (1939, 1940) found in their tumours are considered, this variation in metachromatic properties seems to have been partially explained. In addition, with the great differences in the histology of the various lesions, would be logical to expect certain tumours with marked adenoid appearance to secrete more than those with an essentially non-glandular structure.

Three of the five autopsies performed show evidence of peritoneal metastases. In the other two intra-thoracic spread was observed, in one case to the other lung and chest wall, and in the second to the mediastinal lymph glands and pericardium. Biopsy evidence of metastases has been obtained in two cases, one from the omentum and the other from the subcutaneous tissue of the chest wall; while a third patient developed an implantation nodule in the thoracotomy scar which showed a similar appearance to that of the previous biopsy specimen.

The first recorded case of carcinoma of the lung associated with asbestos was described by Lyall and Smith (1935). By 1955, according to Doll a total of 61 cases had been reported. Included in these cases was one mesothelioma. Cartier (1953) mentioned two cases of diffuse mesothelioma from the Canadian chrysotile mine. A further three cases were described by van der Schoot (1953). Unfortunately no indication is given in the literature regarding the type of asbestos to which the majority of recorded cases of carcinoma were exposed. However, discussion with management and medical officers of two of the factories, in which the majority of the cases reported in Britain were employed, suggests that most of these workers were handling chrysotile asbestos (Wagner, 1953). The possibility that some of these people may also have been exposed to crocidolite dust cannot be excluded.

Attempts to produce tumours in experimental animals by exposing them to asbestos dust have not been made. Vorwald and Karr (1938) were unsuccessful. Lynch, Melver, and Cain (1957) succeeded in producing tumours in mice, but lung neoplasms in control animals were far too numerous for the results to be considered significant. On the other hand Schmähl (1958) working in Druck's laboratory has been able to produce sarcomas in rats. This has occurred after subcutaneous

intrapleural inoculation of both asbestos fibres and dust. He states that "mineral asbestos" was used but does not name the variety.

In our series of mesotheliomata, histological evidence of asbestosis has been observed in eight of the 10 cases in which lung parenchyma was included in the specimen examined. No lung tissue was present in the biopsies from the remaining 23 cases. It was only in the four cases that came to necropsy that large sections of the lung tissue were available for examination. In all of these specimens evidence of asbestosis was found.

At first it was thought that the presence of numerous asbestos bodies, fragments of fibre, and dust immediately below the pleural elastic laminae in these specimens, might have been significant in the pathogenesis of the tumours. However, it is more probable that this distribution is a result of the marked atelectasis. Similar features have been observed in atelectatic lungs of asbestos miners in which no mesotheliomas have been observed.

Pleural fibrosis has been a common finding in cases of asbestosis and in some cases large pleural plaques measuring up to 1.0 cm. in thickness have been seen. These plaques, which were first described by Gloyne (1933), have been observed following exposure to both amosite and crocidolite asbestos dusts. In these cases of benign pleural thickening no evidence of stromal metachromasia has been observed.

In all the histological sections of pleura examined in the cases of these mesotheliomas and in more than 100 cases of asbestosis, no asbestos bodies, fragments of fibres, or dust have been observed beyond the pleural elastic laminae.

Three patients from the Kuruman district, with clinical and radiological features consistent with those of diffuse pleural mesothelioma had markedly abnormal cells in the pleural fluid. These cells showed no specific features to distinguish them from cells originating in secondary malignant pleural deposits nor from the grossly atypical cells sometimes seen in non-malignant pleural effusions. As no biopsy or autopsy examinations were obtained in these cases they have been excluded from the series.

The pathological evidence for associating these tumours with asbestos exposure is not conclusive. As previously stated, only in eight of the 33 cases has evidence of asbestos been demonstrated. Of these, six had a definite mining history and one had been exposed to asbestos while lagging steam pipes. The other case was born in the Kuruman district and nothing else is known of him, until his arrival at a Witwatersrand gold-mine at the age of 23. In the remaining 25 cases we can only present circumstantial evidence of exposure to asbestos dust

(Table 3). Eighteen of these 25 cases were born in the vicinity of the mines and two arrived in the district as infants. Of these 18 people 11 admit definite childhood exposure to the dust and two others were exposed industrially in later life. In addition two patients with childhood exposure later worked in the asbestos mines. Three cases arrived in the region at an older age but were employed either on the mines or in transporting asbestos. A further three of these 25 cases have had industrial exposure, and in only one case do the relatives deny any exposure to asbestos dust.

The four industrial cases are significant. Two of the patients were lagging locomotive boilers and one was lagging steam pipes. A man, who was an upholsterer by trade, was employed in making fire-proof clothing from 1939-1945. As far as can be ascertained these people were never in the Griqualand district. These findings tend to add support to asbestos being the common factor in the development of these tumours, and to counter the suggestion that there may be some other environmental cause in the region of Griqualand West.

If asbestos dust is a factor in the occurrence of these tumours, similar cases might have been expected from the neighbourhood of the Transvaal asbestos mines in the Pietersburg and Lydenburg districts, where crocidolite and amosite asbestos is mined. According to Vermaas (1952), crocidolite and amosite occur in the same seams in the Pietersburg district. As can be seen from Table 1, amosite is similar in composition to crocidolite. In the past four years the lungs of 24 cases of asbestosis from the Lydenburg district have been examined. In this material two cases of adeno-carcinoma have been observed. The one was in a white miner with 19 years' service, who had an adeno-carcinoma arising from a bronchus. The other case was a Bantu miner who had a peripheral tumour. No service record was obtainable in this case. Only one case of asbestosis has been received from the Pietersburg area. All of these men were actually employed on the mines at the time of death, the majority having had a relatively short service. Our findings suggest that mesothelioma occurs 20 to 40 years or more after exposure to dust. Until comparatively recently the mining in the Transvaal has been on a small scale, and there were no settlements in the vicinity of the mines.

The lungs of 20 asbestos miners from the Cape Asbestos Field have been examined, in whom no mesothelioma was observed. One autopsy specimen consisting of three fragments of lung showed asbestosis and an adeno-carcinoma. Radiological features of asbestosis have been observed in many miners from this region, without any suggestion of

TABLE 3  
DIFFUSE PLEURAL MESOTHELIOMA: ASSOCIATION WITH ASBESTOS

(1) Case No.	(2) Year of Birth	(3) Age at Diagnosis	(4) Race	(5) Sex	(6) Born on Asbestos Fields	(7) Asbestos Exposure	(8) Diagnosed on Biopsy	(9) Necropsy	(10) Histological Evidence of Asbestosis	(11) Survival (Years)
1	1920	36	R	M	+	Other history unknown until came to the Witwatersrand at the age of 23	-	-	-	1
2	1913	42	MXD	M	+	Mined asbestos from 1930-33; left area at the age of 27	-	+	+	2
3	1902	53	B	F	+	Lived whole life in a location near an asbestos mill	+	-	-	1
4	1896	58	W	F	+	Lived on asbestos fields until the age of 5; worked in asbestos warehouse 1916-20	+	-	-	20
5	1925	31	R	M	+	Spent all his working life in the vicinity of mines	-	+	-	11
6	1903	53	W	F	+	Lived all her life in the vicinity of mines	+	-	-	15
7	1920	36	W	M	+	Lived all his life in the vicinity of mines; worked as a miner	+	-	+	24
8	1894	63	MXD	F	+	From the age of 24 lived in a village serving local mines; often visited mines; watched cobbing outside houses	+	-	-	5
9	1903	52	MXD	M	+	Whole life spent near mines, digging wells	+	-	-	5
10	1909	49	W	M	+	Lived at the mine from age 7-17 years; played on dumps and in mine as a boy; returned to assist from age 21-23	+	-	-	13
11	1898	60	MXD	M	+	Whole life spent near mines; miner 1931-33	+	+	+	12
12	1910	48	B	M	+	Lived near mines until the age of 17; miner 1927	+	-	+	Still
13	1909	50	W	F	+	Lived near mine until the age of 21; played with fibre as a child	+	-	-	7
14	1918	40	B	M	+	Whole life spent near mines; miner 1938-41; played on dumps as a child	+	+	+	9
15	1916	42	W	F	+	Daughter of Case 22; lived at mine until age of 20; went to school near cobbing sheds	+	-	-	10
16	1896	60	W	M	+	Went to school near mines; transported asbestos 1914-16	+	-	0	11
17	1911	48	MXD	F	+	Lived on major wagon route till age of 15; several subsequent visits	+	-	-	3
18	1920	38	R	M	+	Whole life in the vicinity of the mines; miner 1945-58	+	+	+	12
19	1922	37	MXD	M	+	Family lived at mine; miner 1938-1959	+	-	+	10
20	1906	53	W	F	+	Spent whole life in village on wagon route to Kimberley	+	-	+	8
21	1912	44	W	M	+	Lived in the vicinity of mines until the age of 16; often on dumps as a child	+	-	-	5
22	1889	68	W	M	+	Whole life in the vicinity of the mines; miner 1913-22, and 1945-52	+	-	-	2
23	1893	63	W	F	+	Lived in the vicinity of the mines until the age of 30	+	-	-	12
24	1922	35	W	M	-	Lived in the vicinity of a mill from the age of 1-7; played on the dumps as a child	+	-	-	5
25	1899	50	W	F	-	Lived in a mining area from the age of 10 to 18 years; after 1918 spent whole life in same town as Case 24	+	-	-	10
26	1895	49	W	M	-	Mined and transported asbestos from 1928-33 as overseer	+	-	-	3
27	1904	52	W	M	-	Born in North West Cape; transported asbestos from 1920-24	+	-	0	5
28	1899	60	W	M	-	Lived whole life on farm in mining area from the age of 12; transported asbestos 1916-21	+	-	-	7
29	1913	44	W	M	-	Maintaining locomotive boilers 1931-45	+	-	-	1
30	1909	50	W	M	-	Maintaining steam pipes in explosive factories 1930-40	+	-	+	1
31	1913	44	W	M	0	Worked as fitter on railways, maintaining locomotive boilers, dates unknown	+	-	-	1
32	1908	49	W	M	-	Making asbestos blankets for the Air Force 1939-45	+	-	-	1
33	1890	59	W	F	-	No history of exposure to asbestos	+	-	-	1

KEYS: Column 4—  
Race: W = White  
MXD = Mixed (Coloured)  
B = Black

Column 6  
+ = Born on asbestos fields  
- = Not born on asbestos fields  
0 = Unknown

Column 9—  
+ = Necropsy done  
- = Necropsy not done  
\* = Only small fragments of parietal pleura submitted at autopsy for histological examination

Column 10  
+ = Positive histological evidence of asbestosis  
- = No lung tissue submitted  
0 = Small fragments of lung tissue, but no evidence of asbestosis

tumour formation. However, pleural involvement has been a common finding in these cases (Hurwitz, 1959). It is possible that there has not been a sufficient lapse of time after exposure for tumour development in these cases. Further, the factor

of individual susceptibility must also be considered.

We wish to record the great assistance that we have received in this investigation from the medical practitioners of the Griqualand West District and the

and pathologists in Johannesburg, Durban, and Durban.

I thank the Secretary of Health for permission to publish this paper.

## REFERENCES

11. G. and Bova, G. (1937). *Acta med. port.*, 17, 367.  
 Hill, W. H. (1950). *Amer. J. Path.*, 26, 471.  
 H. P. (1952). *Arch. industr. Hyg.*, 5, 262. (Contribution to the discussion.)  
 R. (1955). *Brit. J. industr. Med.*, 12, 81.  
 J. G. E. D. (1955). *Memorandum on the Asbestos Industry in the Cape Province*. Report of the Government Mining Engineers, Union of South Africa, pp. 76-82.  
 M. S. R. (1931). *Tubercle (Lond.)*, 14, 493.  
 G. M. G. (1957). *Cancer (Philad.)*, 10, 298.  
 C. W. (1946). *Nature (Lond.)*, 157, 802.  
 A. L. (1930). *Asbestos in the Union of South Africa*. Memorandum no. 12. Geological Survey of South Africa.  
 G. J. S. (1959). Personal communication.  
 G. J. S. and Oettle, A. G. (1957). *Acta Un. int. Cancer.*, 13, 949.  
 H. M. (1959). *Proceedings of International Conference of Experts on Pneumoconiosis*, Johannesburg, Feb. 1959. Churchill, London. (In the press.)  
 H. L. E. (1947). *Amer. J. Path.*, 23, 871.  
 H. P. and Kahn, C. B. (1931). *Arch. Path. (Chicago)*, 11, 335.  
 H. L. (1953). *Histochemie et Cytochemie Animales*, pp. 332-337. Gauthier-Villars, Paris.  
 H. N. M. and Smith, W. A. (1935). *Amer. J. Cancer*, 24, 56.  
 Melzer, F. A., and Cain, J. R. (1957). *A.M.A. Arch. industr. Hyg.*, 15, 207.  
 Martiny, O. (1956). *Proc. Trans. Mine med. Offrs' Ass.*, 35, 63.  
 Maximow, A. (1927). *Arch. exp. Zellforsch.*, 4, 1.  
 McCaughey, W. T. (1938). *J. Path. Bact.*, 76, 317.  
 Meyer, K., and Chaffin, E. (1939). *Proc. Soc. exp. Biol. (N.Y.)*, 42, 797.  
 — (1940). *J. Biol. Chem.*, 133, 83.  
 Novak, E. (1931). *Amer. J. Obstet. Gynec.*, 22, 826.  
 Robertson, H. E. (1924). *J. Cancer Res.*, 6, 317.  
 Sano, H., F. Weiss, E., and Gault, E. S. (1950). *J. thorac. Surg.*, 19, 781.  
 Schnuhl, D. (1958). *Z. Krebsforsch.*, 62, 561.  
 Smart, J., and Hinson, K. F. W. (1957). *Brit. J. Tuberc.*, 51, 319.  
 Stout, A. P., and Murray, M. R. (1942). *Arch. Path. (Chicago)*, 34, 951.  
 Tobiassen, G. (1955). *Acta path. microbiol. scand.*, Suppl. 105, p. 198.  
 van der Schuer, H. C. M. (1958). *Ned. T. Geneesk.*, 102, 1124.  
 Vermeas, F. H. S. (1952). *Trans. geol. Soc. S. Afr.*, 55, 199.  
 Vorwald, A. J., and Karr, J. W. (1938). *Amer. J. Path.*, 14, 49.  
 Wagner, J. C. (1958). *Memorandum on Pneumoconiosis Research in Europe*. Submitted to the C.S.I.R.  
 Willis, R. A. (1948, 1st ed. and 1953). *Pathology of Tumours*. Butterworth, London.

## Addendum

By the end of June 1960, a total of 47 cases of mesothelioma had been identified. In 45 of these a possible association with exposure to crocidolite has been established. In one case a mesothelioma of the peritoneum was present.

## THE JULY (1960) ISSUE

The July (1960) issue contains the following papers:—

- Compression Sickness during Construction of the Dartford Tunnel. By F. Campbell Golding, P. Griffiths, H. V. Hempleman, W. D. M. Paton, and D. N. Walder.
- Comparison of the "High-altitude" and "High-pressure" Syndromes of Decompression Sickness. By M. de G. Gribble.
- Asphyxiant Poisoning Caused by the Use of Chemical Paint Removers Containing Methylene Chloride in Ill-ventilated Rooms Heated by Kerosene Stoves. By W. B. Gerritsen and C. H. Buschmann.
- Study on the Acute Toxicity of the Tri-aryl Phosphates Used as Plasticizers. By H. F. Bondy, E. J. Field, A. N. Worden, and J. P. W. Hughes.
- Lead Intoxication of CaNa<sub>2</sub>EDTA (Moxatil) by Workers Constantly Exposed to Lead Poisoning. By Lj. Petrović, M. Stanković, M. Savičević, and D. Poletti.
- Biochemical Study of the Urinary Protein of Men Exposed to Metallic Mercury. By J. C. Smith and Agnes R. Wells.
- Simplified Method for the Estimation of Nickel in Urine. By J. Gwynne Morgan.
- Physical, Bronchographic, Radiological, and Physiological Observations in Ten Cases of Asbestosis. By G. L. Leathart.
- Measurement of Occupational Mortality. F. D. K. Liddell.
- Observations on the Occupational Life History of the Coal Face Worker at Two Collieries. By O. P. Edmonds and D. S. Kerr.
- Assessment of Electroencephalographic Changes and Memory Disturbances in Acute Intoxications with Industrial Poisons. By B. Chalupa, J. Synková, and M. Ševčík.
- Book Reviews.

A number of copies are still available and may be obtained from the Publishing Manager, British Medical Association, Tavistock Square, W.C.1, price 17s. 6d.

Article by H.C.H. VAN DER SCHOOFF: "ASBESTOSIS AND PLEURAL TUMORS" June 7, 1933  
NETHERLANDS JOURNAL FOR MEDICINE, VOL: 192, ( 1 ) 23, p 1125-1126

" Since Lynch and Smith, in 1935, published the first case of lung cancer in a victim of asbestosis, numerous communications have appeared in the literature on the concurrence of both processes. The carcinomas thus described appear generally to have had their origin in the peripheral branches of the bronchus and could be classified histologically as epidermoid carcinoma, oatcell adenocarcinoma or as anaplastic carcinoma, with epidermoid carcinoma predominating in this series. In the literature only one case is mentioned of the combination of asbestosis and pleural tumor, without an account of the localization or histological detail. During the past year we encountered this combination ( of asbestosis and pleural tumor ) twice and to which we now can, through the courtesy of Dr SCHONWAGEL, add a third case.

In all three medical histories we are concerned with workers who were for a protracted period engaged in the occupation of " insulator ". Their work consisted in part disassembling old insulation, which involved often performance of work in poorly ventilated spaces in which a great amount of dust, with a high concentration of asbestos fibers, was released.

<sup>A</sup> Patient A was a man of 67 years who had worked for 41 years with asbestos. One year before his death he was admitted for a hemorrhagic pleuritis (rightsided). No tumor cells were detected in repeated aspirates of the exudate. After two months, the manifestation of pleuritis disappeared.

Eight months later the patient was readmitted in moderately ill condition, with severe dyspnea. On fluoroscopy a considerable pleural thickening was seen on the right side together with a lung picture which suggested asbestosis. No asbestos bodies were however found in the sputum. While the patient was being studied he developed a bronchopneumonia, from which he died.

At autopsy a very large tumor was found which enveloped the whole lung and which had infiltrated through the diaphragm. On microscopic examination numerous asbestos bodies were found in both lungs, with only slight fibrosis of the lung tissue. The tumor showed the characteristics of a mesothelioma and in all probability had arisen from the pleura.

Patient B also was an insulator aged 67 years, who had worked with asbestos for 13 years, and who had experienced symptoms of cough and shortness of breath for 8 years prior to his death. After these symptoms had endured for five years a diagnosis of pulmonary asbestosis was made on the basis of the medical history and the X-Ray picture and he was determined to be 100 per cent incapacitated.

Half a year later a left-sided pleural exudate appeared, in which no tumor cells were detected. This exudate disappeared later, but on the X-ray (chest) increasing opacification of the lower left lung field was observed. Two years later the patient died with the status of increasing respiratory failure.

At necropsy the pleura of both lungs was found to be markedly thickened, and a tumor mass, which had enveloped the lower lobe of the left lung and which had infiltrated the diaphragm, was found. Microscopic analysis verified the presence of asbestosis which was of quite marked degree. The histological characters of the tumor was more or less identical with that of patient

aged

Patient C (Dr Schormann's) was an insulator 57 years for whom it was not determined how long he had worked in the insulation industry. His history is shorter and shows little correspondence with that of the other two cases. Half a year before his demise the patient complained of severe pain in the upper part of abdominal region, for which no organic abnormality was initially detected. There were no respiratory system complaints. During his stay in the hospital, however, ossification of the lower portion of the right lung field arose and a tumor became palpable below the right costal margin. A little fluid was aspirated from the right pleural cavity in which malignant neoplastic cells of undiagnosable origin were demonstrable. The patient expired after several months in a state of cachexia.

At autopsy a fibrosarcoma was found. It had surrounded the lower and middle lung lobes and had grown through the diaphragm deep into the liver. Only a moderate number of asbestos bodies were present in the lungs.

bronchus

A long series of anecdotal reports on the concurrence of cancer and asbestosis has finally lead to the universal acceptance of the opinion (point of view) that asbestos should be regarded as one of the factors in the genesis of lung cancer. Asbestos, which promoting / etiological consists of magnesium silicate, does not contain any carcinogenic material, so that the carcinogenic action of asbestos fibers must be ascribed to the chronic mechanical irritation. In conformance herewith, the lung-cancer tends to develop in lung regions where the asbestos fibers occur in greatest concentration, in general thus in the lower lung lobes, particularly in the vicinity of the diaphragm and the lung periphery where the asbestos fibers exert their strongest irritation because the greatest ventilatory excursions occur here.

From the fact that in pulmonary asbestosis pleural thickening, including plaque formation, seldom is lacking, one can assume that mechanical irritation by the asbestos fibers contributes at the very least to the thickening of the pleural layers.

lumen of the

In the above described cases no asbestos fibers were detected in the bronchi or bronchioles in any of the numerous sections. Metaplastic deviation of the bronchial epithelium, such as that which had been demonstrated so clearly by HANKE in 1952 in a patient who suffered from asbestosis and lung carcinoma, also was lacking. If one assumes that the dimensions of inhaled asbestos fibers are of such a size that they can pass unimpeded through the respiratory tree so that their irritating action would take place in the peripheral portions of the lungs, it is comprehensible why pleural thickening dominates the picture.

These considerations lead to drawing the line of correlation between lung cancer and asbestosis at the level of the pleural thickening. We are fortified in this by the investigations of DOLL, published in 1955, in which he showed on statistical grounds that the average chance of lung cancer in asbestos workers, who have practiced their profession for 20 or more years, is ten times as great as compared with the population in general. His table, which only mentioned the histological diagnosis, included one case of pleural endothelioma, and he thus mentions this pleural tumor in the same breath with lung tumors.

According to the data of the central service of the labor service, asbestosis occurs by far most frequently in the west of our country where the insulation industry is established. Here there are hundreds of workers. Precise data concerning the prevalence of asbestosis are not presently available, but it can be generally assumed that about 20 per cent of the workers suffer in

lesser or greater degree from asbestosis. Prevention of asbestosis appears to be a difficult technical <sup>or</sup> problem.

DISCUSSION:

Dr VAN DER ZALM: drew attention to a case of mesothelioma of the pleural cupola which was detected next to silicosis with pseudumor formation. Are there in the literature more similar cases?

DR VAN DER SCHOOT: ( response ) : The literature which I have reviewed has not revealed any publication which establishes a link between pleural tumors and any form of pneumoconiosis. The relationship between silicosis and cancer is decidedly less clear.

DrSCHONWAGEL: Recently I performed an autopsy on yet another deceased patient with asbestosis, who, however, had a mesothelioma of the peritoneum. Would you be able to establish a connection between this asbestosis and the peritoneal tumor? The patient was an insulator and worked temporarily in an atomic factory in Australia.

DR VAN DER SCHOOT ( Answer ) : If it is established that there was no tumor tissue within the thorax and that the tumor had not grown through the diaphragm, it would seem to me difficult to establish a connection between the two disease processes.